

Twin Research and Human Genetics

<http://journals.cambridge.org/THG>



Additional services for *Twin Research and Human Genetics*:

Email alerts: [Click here](#)

Subscriptions: [Click here](#)

Commercial reprints: [Click here](#)

Terms of use : [Click here](#)

The UK Adult Twin Registry (TwinsUK Resource)

Alireza Moayyeri, Christopher J. Hammond, Deborah J. Hart and Timothy D. Spector

Twin Research and Human Genetics / *FirstView* Article / November 2012, pp 1 - 6

DOI: 10.1017/thg.2012.89, Published online: 22 October 2012

Link to this article: http://journals.cambridge.org/abstract_S1832427412000898

How to cite this article:

Alireza Moayyeri, Christopher J. Hammond, Deborah J. Hart and Timothy D. Spector The UK Adult Twin Registry (TwinsUK Resource). *Twin Research and Human Genetics*, Available on CJO 2012 doi:10.1017/thg.2012.89

Request Permissions : [Click here](#)

The UK Adult Twin Registry (TwinsUK Resource)

Alireza Moayyeri, Christopher J. Hammond, Deborah J. Hart, and Timothy D. Spector

Department of Twin Research and Genetic Epidemiology, King's College London, St. Thomas' Hospital, London, UK

TwinsUK is a nation-wide registry of volunteer twins in the United Kingdom, with about 12,000 registered twins (83% female, equal number of monozygotic and dizygotic twins, predominantly middle-aged and older). Over the last 20 years, questionnaire and blood/urine/tissue samples have been collected on over 7,000 subjects, as well as three comprehensive phenotyping assessments in the clinical facilities of the Department of Twin Research and Genetic Epidemiology, King's College London. The primary focus of study has been the genetic basis of healthy aging process and complex diseases, including cardiovascular, metabolic, musculoskeletal, and ophthalmologic disorders. Alongside the detailed clinical, biochemical, behavioral, and socio-economic characterization of the study population, the major strength of TwinsUK is availability of several 'omics' technologies for the participants. These include genome-wide scans of single nucleotide variants, next-generation sequencing, exome sequencing, epigenetic markers (MeDIP sequencing), gene expression arrays and RNA sequencing, telomere length measures, metabolomic profiles, and gut flora microbiomics. The scientific community now can freely access parts of the phenotype data from the 'TwinsUK', and interested researchers are encouraged to contact us via our Web site (www.twinsuk.ac.uk) for future collaborations.

■ **Keywords:** twin studies, genetic epidemiology, longitudinal studies, genome-wide association studies, next-generation sequencing, genomics, epigenomics, gene expression analysis, metabolomics, microbiomics

Introduction

The UK Adult Twin Registry (TwinsUK) is a cohort of volunteer adult twins from all over the United Kingdom. The Department of Twin Research and Genetic Epidemiology at St. Thomas' Hospital, King's College London (KCL) hosts the registry, which started in 1992 via media campaigns targeted at middle-aged women. The success of early studies led to rapid evolution of the registry and it now incorporates twins, both male and female, from other sources such as the Aberdeen Twin Registry and Institute of Psychiatry Adult Registry. The primary focus of study has been the genetic basis of complex diseases (cardiovascular, metabolic, musculoskeletal, and ophthalmologic diseases), which has broadened to include the complex healthy ageing process. The third health check of the volunteer twins has provided longitudinal data that, alongside with the state-of-the-art 'omics' technologies data, can significantly advance the field of genetic and clinical epidemiology of ageing. We have previously described the design of the original twin registry, facilities and procedures for data collection, clinical and biological assessments, and main findings in 2006 (Spector & Williams, 2006). The current article presents a brief update on the study procedures and recent technological

applications in our cohort. More detailed description of phenotypes, research projects and collaborations, papers, and study findings can be accessed through our updated study Web site (<http://www.twinsuk.ac.uk>).

The Collection

The TwinsUK registry now consists of about 12,000 monozygotic (MZ) and dizygotic (DZ) twins aged 18 to 103 years (Table 1). About 83% of the registry is female (mean age of 55 years). The registry now contains 51% MZ and 49% DZ twins. Between 1992 and 2004, twins were invited for a full comprehensive visit and several project-led studies. More than 7,000 twins responded to some of the annual questionnaires and 5,725 attended a comprehensive visit. Apart from a lifelong lower weight in MZ twins of around 1 kg, all other age-matched characteristics of these

RECEIVED 1 September 2012; ACCEPTED 16 September 2012.

ADDRESS FOR CORRESPONDENCE: Professor Timothy D. Spector, Department of Twin Research and Genetic Epidemiology, King's College London, St. Thomas' Hospital, London SE1 7EH, United Kingdom. E-mail: tim.spector@kcl.ac.uk

TABLE 1
The UK Adult Twin Registry Update

Name of register	UK Adult Twin Registry (TwinsUK)
Country	United Kingdom
Kind of ascertainment	Volunteers unselected
Opposite-sex twins (yes or no)	Yes
Number of pairs	1900–1920: 22 FF; 2 MM; 2 FM
(Separated by birth range and sex)	1920–1930: 390 FF; 70 MM; 4 FM 1930–1940: 1,518 FF; 189 MM; 32 FM 1940–1950: 2,480 FF; 308 MM; 58 FM 1950–1960: 2,172 FF; 429 MM; 88 FM 1960–1970: 1,889 FF; 486 MM; 62 FM 1970–1980: 1,356 FF; 394 MM; 42 FM 1980–2000: 558 FF; 124 MM; 40 FM
Grand total	10,393 FF; 2,002 MM; 328 FM (6,369 pairs)
Major interests	Common complex diseases and ageing traits
Traits measured	Full questionnaires and clinical examinations on majority of twins for wide range of over 1,000 clinical and biochemical traits including: cardiovascular diseases, obesity, metabolic syndrome, respiratory diseases, dermatology, osteoarthritis, osteoporosis, eye diseases, back diseases, coagulation system, immune function, cognitive function, gastro-intestinal system, pain thresholds, allergy, atopy, sexuality, pitch perception, and various aspects of personality.
DNA samples	13,458 aliquots from 7,548 twins 9,321 aliquots from 5,965 twins stored as back-up DNA samples from 995 parents and 1,227 siblings taken and stored
Other samples	119,511 blood samples (65,980 serum, 43,527 plasma EDTA, 10004 plasma Li heparin) from 7,681 twins (16,677 back-up samples) 28,276 urine samples from 7681 twins (16,677 back-up samples) Range of 2–76 aliquots of various specimens at multiple time points
Comments	Monozygotic:Dizygotic ratio is approximately 1:1 Majority of twins are female with mean age of 55 years 5,710 twins with genome-wide association data 5,000 twins with DNA methylation data by the end of 2012 2,000 twins with next-generation sequencing data by the end of 2012 Data available for transcriptome across multiple tissues, telomere length, and metabolomic profile in different subsamples
Main sources of funding	Wellcome Trust, UK Medical Research Council (MRC), British Heart Foundation, NIHR Biomedical Research Centre, Pfizer, Chronic Disease Research Foundation, and the European Union framework programs
Contact	Prof. Timothy D. Spector
Address	Department of Twin Research and Genetic Epidemiology, St. Thomas' Hospital, London SE1 7EH
E-mail	tim.spector@kcl.ac.uk or victoria.vazquez@kcl.ac.uk
Web site	www.twinsuk.ac.uk

Note: FF = female–female, MM = male–male, FM = female–male.

volunteer twins were found not to differ from a singleton population-based cohort of British women (Chingford study; Andrew et al., 2001). Between April 2004 and May 2007, all the 6,740 active twins on the registry were invited for a 1-day clinical visit, of whom 3,725 twins attended and 1,299 twins posted their blood DNA samples via their general practitioners. The age of participants ranged between 18 and 82 years (mean 52.5 ± 13 years) and 3,299 of the clinic attendants (89%) were female.

The second follow-up visit, also known as the HATS (Healthy Ageing Twin Study) visit, started in August 2007. Only women aged ≥ 40 years with at least one previous clinical visit ($n = 4,610$) were invited for this visit. In total, 3,125 women (mean age 59.6 ± 9 years, 48% MZ twins) attended the clinic (response rate = 68%). Follow-up time between first and last visits ranged between 6.1 and 17.4 years (mean 11.2 ± 2 years). Six hundred of the participants in this visit had four or more previous clinical visits. Compared to the baseline and first follow-up visits, participants in the HATS visit appeared to have higher socio-economic status, lower self-rated health status, and be more health aware given their level of alcohol intake and smoking. Comparison of the respondents and non-respondents to the HATS

invitation showed that the non-respondents were generally younger and of lower socio-economic status. However, no significant clinical differences were observed between attendants and non-attendants to the HATS visit (Moayyeri et al., 2012b), suggesting that the participants in this visit can be considered as representative of the original population in the study. Zygosity status was assessed for all twins at the time of registration by a 'peas in a pod' questionnaire and confirmed via subsequent genotyping or genome-wide association studies.

Longitudinal Data

Longitudinal data is available for a wide range of phenotypes based on different questionnaires, clinical tests, and biochemical assays performed at different stages of the study (Moayyeri et al., 2012b). Table 2 summarizes some of these clinical measures showing the breadth of the available data and the potential for studies on various aspects of healthy ageing in this population. For instance, longitudinal changes in bone mineral density in more than 4,000 twins over an average of 17.5 years serves as a valuable endpoint for clinical and genetic epidemiological studies (Moayyeri

TABLE 2
Longitudinal Data Available in the TwinsUK Registry Participants

Phenotype	N of measurements	N of participants	N of participants with ≥ 2 measurements	Maximum number of visits	Duration of follow-up (year)	Maximum duration of follow-up (year)
Blood pressure	13,180	7,189	3,836	5	7.3 \pm 3.3	13.7
Lipid profile	12,652	6,881	3,585	7	8.5 \pm 3.9	16.4
Electrocardiography	8,083	5,533	1,864	5	8.0 \pm 3.1	12.6
Fasting glucose	16,305	7,731	4,344	11	8.4 \pm 3.6	16.8
Blood insulin	13,650	6,953	3,810	11	8.4 \pm 3.6	16.8
Respiratory function	12,245	7,128	3,326	5	8.7 \pm 3.5	16.2
Hip BMD	15,367	7,025	4,098	12	8.5 \pm 3.8	17.5
Spine BMD	15,489	7,046	4,115	13	8.5 \pm 3.8	17.5
Whole body DXA	13,850	7,292	3,882	7	8.5 \pm 3.7	17.5
Heel QUS	8,299	5,419	1,753	7	3.4 \pm 1.1	7.7
Grip strength	6,861	4,840	1,999	3	3.1 \pm 0.7	5.5
Sexual hormones	5,779	5,350	428	3	5.3 \pm 3.0	10.3

Note: BMD = bone mineral density, DXA = dual-energy X-ray absorptiometry, QUS = quantitative ultrasound.

et al., 2012a). During the study follow-up, care has been taken to perform key clinical tests with similar protocols across all visits. Incident clinical endpoints (e.g., cardiovascular events, stroke, fractures, osteoarthritis, and different cancers) have been assessed over the course of study using questionnaires. Twins are all now registered using their National Health Services (NHS) numbers with the Office for National Statistics of England for retrospective analysis and future follow-up regarding their cancer and mortality status.

Novel Molecular and Genetic Phenotypes

Alongside with the conventional epidemiological phenotypes assessed by questionnaires and clinical visits, the TwinsUK registry benefits from generous and continued donation of biological samples by its volunteering participants. The methods for collection of these biological samples have been described previously, and the updated figures are presented in Table 1. Recently, a wide array of the latest 'omics' advances has been applied to subsections of these samples that makes TwinsUK one of the most uniquely phenotyped and deeply genotyped populations in the world. Here we describe some of these advances.

Genome-Wide Association Studies

TwinsUK has contributed to many international consortia for genome-wide association analysis of various phenotypes. Genome-wide scan data using two chips (Illumina HumanHap300 BeadChip and Illumina HumanHap610 QuadChip) are available for 5,710 twins. The data have been fully imputed using 'HapMap II' and '1000 Genomes' reference panels (containing ~ 2.5 and ~ 16 million single nucleotide polymorphisms, respectively). TwinsUK is a member of many ongoing international consortia for meta-analysis of various traits, such as GIANT, CHARGE, ENGAGE, GEFOS, SUNLIGHT, MolPAGE, VisiGEN, TreatOA, and SpiroMeta (Aulchenko et al., 2009; Dehghan et al., 2011; Duffy et al., 2010; Dupuis et al., 2010; Elks et al., 2010;

Evangelou et al., 2011; Ganesh et al., 2009; Hysi et al., 2010; Kolz et al., 2009; Lango et al., 2010; Lindgren et al., 2009; Newton-Cheh et al., 2009; Nolte et al., 2009; Padmanabhan et al., 2010; Panoutsopoulou et al., 2011; Repapi et al., 2010; Richards et al., 2009a, 2009b; Smith et al., 2010; Speliotes et al., 2010; Willer et al., 2009; Zhai et al., 2009, 2011).

Next-Generation Sequencing

The UK10K study is an ongoing collaboration between TwinsUK study, the Wellcome Trust Sanger Institute, and several other collaborators for using the state-of-the-art, next-generation sequencing methods to uncover rare genetic variants associated with health and disease. The study involves whole-genome sequencing of 4,000 healthy people with well-documented physical characteristics (2,000 twins from TwinsUK and 2,000 children from ALSPAC study) and whole-exome (protein-coding regions of DNA) sequencing of 6,000 people with extreme health problems (obesity, neurological problems, and rare diseases). At present, all twin samples have been sequenced (6 \times depth) and both phenotype and sequence data will be publicly available soon. More details about the study can be accessed at: www.uk10k.org. Moreover, about 1,000 exome sequences at 30–60 \times depth have been performed for twin participants as part of projects with Pfizer and the GoT2D consortium. Over 2,000 Exome chips are currently being performed, mainly for control purposes in other consortia.

Epigenetic Markers

The first epigenetic assessment in TwinsUK was performed on DNA methylation patterns using Illumina HumanMethylation27 BeadChip in a sample of 172 female twins. This array examines 27,578 promoter CpG-sites that map uniquely across the genome, and some of these sites were found to be associated with age and age-related phenotypes (Bell et al., 2012). Currently, the Infinium HumanMethylation450 BeadChip (Illumina) is being applied to 500 additional MZ and DZ twin pairs to generate

higher-resolution genome-wide DNA methylation profiles. This array includes 485,764 cytosine positions (CpG dinucleotides and CNG sites) across the human genome. Meanwhile, the major ongoing epigenetic project using the TwinsUK population is the EpiTwin study (<http://www.epitwin.eu>), which uses MeDIP (Methylated DNA immunoprecipitation) sequencing in whole blood samples (Bell & Spector, 2011). This is the largest epigenetic project of its kind, in collaboration with the Beijing Genomics Institute, aiming to assay epigenomic differences in 5,000 adult UK twins aged 16–85 years, discordant and concordant for a wide variety of diseases and environments. Next-generation sequencing has the potential to prove powerful in detecting disease-related methylation differences at a high level of resolution in a sample of this size. The initial targets of the study include obesity, diabetes, allergy, heart disease, osteoporosis, depression, and longevity, but the method can be applied to every common trait or disease.

Gene Expression Measures

Eight hundred fifty-six twins with detailed clinical profiles have been biopsied during the HATS clinical visit. This has been done in the context of the MuTHER (Multiple Tissue Human Expression Resource) project, which is a Wellcome Trust funded study designed to understand the mechanisms involved in common trait susceptibility via gene expression across multiple tissues (Nica et al., 2011). Gene expression in three tissues of skin, fat, and lymphoblastoid cell lines have been measured using Illumina's whole genome expression array (HumanHT-12 version 3) containing 48,803 probes in three technical replicates. Results for expression quantitative trait loci analysis in 856 twins are freely available in the Web site (<http://www.muther.ac.uk>), and the main paper has recently been published (Grundberg et al., 2012). All of these tissues are now being ribonucleic acid (RNA) sequenced as part of the EuroBATS project (Biomarkers of Ageing using whole Transcriptome Sequencing), which is a 3-year European (EU-FP7) project started in January 2011. EuroBATS aims to discover novel biomarkers of ageing by incorporating novel RNA sequencing, telomere measurement, and bioinformatics techniques (<http://www.eurobats.eu>). Sequencing is being performed using Illumina GAIIx platform and the standard protocol of RNAseq (expecting 10–20 million reads per sample).

Telomere Length

Telomere length, as a marker of cellular senescence and subsequent cell death, was first measured in 3,256 twins with available genome-wide scans. These measures were derived from the mean of the terminal restriction fragment length by using the Southern blot method on DNA extracted from peripheral leukocytes. This data has contributed to detection of several genes implicated to affect biological age (Codd et al., 2010; Mangino et al., 2009). Recently, a larger sample (4,899 twins aged 16–99 years)

has been assessed for telomere length using an established and validated quantitative polymerase chain reaction technique. Quality control has now been finalized and the data is available for collaborations.

Metabolomic Profiles

In 2009, fasting serum concentrations of 163 metabolites were measured for 1,270 twins using electrospray ionization tandem mass spectrometry (Biocrates AbsoluteIDQ technology). This targeted panel of metabolites covers a wide range of known lipids, amino acids, sugars, acylcarnitines, and phospholipids. This data has been used in several outstanding studies (Zhai et al., 2010). More recently, a larger sample of 6,055 twins has been assessed using a new method of non-targeted metabolomic analysis. This new platform (Metabolon Inc., Durham, USA) incorporates two separate ultra-high performance liquid chromatography/tandem mass spectrometry injections (optimized for basic and acidic species) and one gas chromatography/mass spectrometry injection per sample. The platform has detected and quantified concentration of 510 small molecules (299 known and 211 unknown molecules) including amino-acids, lipids, carbohydrates, vitamins, nucleotides, peptides, xenobiotics, and steroids. Genome-wide association studies of ~37,000 traits from 60 biochemical pathways in a subsample of this population has identified several genes involved in metabolic individuality in humans and promises significant advances in future functional studies (Suhre et al., 2011).

Microbiomics

We have recently started a collaborative National Institutes of Health-funded study with Cornell University, aiming to collect gut flora DNA for analysis with 16S sequencing technology in 5,000 twins. In addition, pilot data assaying microbiome diversity from other human body sites, such as skin, oral, and nasal cavities, as well as gut flora metagenomics are currently underway.

Future Directions and Collaborations

Frequent data collection with detailed clinical, biochemical, behavioral, and socio-economic characterization of participants for about two decades provides the opportunity to look at the prospective single-measure and repeated-measure associations for complex diseases and domains of healthy ageing in TwinsUK population. This also offers a unique opportunity to explore personalized medicine. Data collection, database management, biological sample storage, and statistical quality control have been carried out to a high standard. Blood, urine, and DNA sample aliquots from all visits are available for future measurements. We currently use online questionnaires and are actively engaging with our twin participants via e-mail and social networking Web sites. Our 'Volunteer Advisory Panel' helps informed

decisions about the ethics, practicalities, and appropriateness of potential studies.

The TwinsUK registry has a history of numerous successful scientific collaborations, and we remain committed to providing the scientific community with access to the phenotype data from the 'TwinsUK Resource'. A recent Biomedical Resource Grant from the Wellcome Trust is continuing to fund the core functions of TwinsUK. This will enable access to our publicly funded data from the wider scientific community (the 'resource'), which will be separate from the individual research projects performed by academics within the Department of Twin Research at KCL. The TwinsUK Resource is opening up access of data, currently harmonizing and standardizing phenotypic data collected over the last 20 years; a subset of the data is already available for full open access via our Web site (<http://www.twinsuk.ac.uk/data-access>) and a search engine for the available phenotypes has been provided. We have an access committee, which meets weekly and reviews about 20 requests a month (www.twinsuk.ac.uk/data-access/submission-procedure). Researchers are encouraged to find out if TwinsUK resource can help them answer their research questions and get in contact for future collaborations.

Acknowledgments

The authors acknowledge financial support from the Wellcome Trust, the Department of Health via the National Institute for Health Research (NIHR) comprehensive Biomedical Research Centre award to Guy's and St Thomas' NHS Foundation Trust in partnership with King's College London/Arthritis Research Campaign/EC FP6 Programme Grant-512066 (LSHG-CT-2004); MOLPAGE/EC Framework 7 Programme Grant 200800; Treat OA/EC Framework 7 Health-2007-A; ENGAGE/EC Framework 7 Health-2007-2.4.5-4; GEFOS/EC FP6 MRTN-CT-2006-034021; MyEuropa Research Training Network/Chronic Disease Research Foundation (CDRF)/Pfizer Pharmaceuticals/National Health and Medical Research Council (NHMRC)/National Institute of Aging (NIA)/Guide Dogs for the blind Association(GDBA)/Biotechnology and Biological Sciences Research Council (BBSRC). The authors acknowledge the funding and support of the National Eye Institute via an NIH/CIDR genotyping project (R01EY018246-01-1 PI: Terri Young). The authors would also like to thank all the twins who participated and supported this cohort and staff in the Department of Twin Research and Genetic Epidemiology, St. Thomas' Hospital, King's College London.

References

- Andrew, T., Hart, D. J., Snieder, H., de, L. M., Spector, T. D., & MacGregor, A. J. (2001). Are twins and singletons comparable? A study of disease-related and lifestyle characteristics in adult women. *Twin Research*, 4, 464–477.
- Aulchenko, Y. S., Ripatti, S., Lindqvist, I., et al. (2009). Loci influencing lipid levels and coronary heart disease risk in 16 European population cohorts. *Nature Genetics*, 41, 47–55.
- Bell, J. T., & Spector, T. D. (2011). A twin approach to unraveling epigenetics. *Trends in Genetics*, 27, 116–125.
- Bell, J. T., Tsai, P. C., Yang, T. P., Pidsley, R., Nisbet, J., Glass, D., & Deloukas, P. (2012). Epigenome-wide scans identify differentially methylated regions for age and age-related phenotypes in a healthy ageing population. *PLoS Genetics*, 8, e1002629.
- Codd, V., Mangino, M., van der Harst, P., Braund, P. S., Kaiser, M., Beveridge, A. J., & Samani, N. J. (2010). Common variants near TERC are associated with mean telomere length. *Nature Genetics*, 42, 197–199.
- Dehghan, A., Dupuis, J., Barbalic, M., Bis, J. C., Eiriksdottir, G., Lu, C., Pelliikka, N., & Chasman, D. I. (2011). Meta-analysis of genome-wide association studies in >80 000 subjects identifies multiple loci for C-reactive protein levels. *Circulation*, 123, 731–738.
- Duffy, D. L., Iles, M. M., Glass, D., Zhu, G., Barrett, J. H., Höiö, V., & Montgomery, G. W. (2010). IRF4 variants have age-specific effects on nevus count and predispose to melanoma. *American Journal of Human Genetics*, 87, 6–16.
- Dupuis, J., Langenberg, C., Prokopenko, I., Saxena, R., Soranzo, N., Jackson, A. U., . . . Barroso, I., for the MAGIC investigators. (2010). New genetic loci implicated in fasting glucose homeostasis and their impact on type 2 diabetes risk. *Nature Genetics*, 42, 105–116.
- Elks, C. E., Perry, J. R., Sulem, P., Chasman, D. I., Franceschini, N., He, C., & Murray, A. (2010). Thirty new loci for age at menarche identified by a meta-analysis of genome-wide association studies. *Nature Genetics*, 42, 1077–1085.
- Evangelou, E., Valdes, A. M., Kerkhof, H. J., et al. (2011). Meta-analysis of genome-wide association studies confirms a susceptibility locus for knee osteoarthritis on chromosome 7q22. *Annals of the Rheumatic Diseases*, 70, 349–355.
- Ganesh, S. K., Zaki, N. A., van Rooij, F. J., et al. (2009). Multiple loci influence erythrocyte phenotypes in the CHARGE Consortium. *Nature Genetics*, 41, 1191–1198.
- Grundberg, E., Small, K. S., Hedman, A. K., Nica, A. C., Buil, A., Keildson, S., . . . Spector, T. D. (2012). Mapping cis- and trans-regulatory effects across multiple tissues in twins. *Nat. Genet*, 44, 1084–1089.
- Hysi, P. G., Young, T. L., Mackey, D. A., et al. (2010). A genome-wide association study for myopia and refractive error identifies a susceptibility locus at 15q25. *Nature Genetics*, 42, 902–905.
- Kolz, M., Johnson, T., Sanna, S., et al. (2009). Meta-analysis of 28,141 individuals identifies common variants within five new loci that influence uric acid concentrations. *PLoS Genetics*, 5, e1000504.
- Lango, A. H., Estrada, K., Lettre, G., et al. (2010). Hundreds of variants clustered in genomic loci and biological pathways affect human height. *Nature*, 467, 832–838.
- Lindgren, C. M., Heid, I. M., Randall, J. C., et al. (2009). Genome-wide association scan meta-analysis identifies

- three Loci influencing adiposity and fat distribution. *PLoS Genetics*, 5, e1000508.
- Mangino, M., Richards, J. B., Soranzo, N., et al. (2009). A genome-wide association study identifies a novel locus on chromosome 18q12.2 influencing white cell telomere length. *Journal of Medical Genetics*, 46, 451–454.
- Moayyeri, A., Hammond, C. J., Hart, D. J., & Spector, T. D. (2012a). Effects of age on genetic influence on bone loss over 17 years in women: the Healthy Ageing Twin Study (HATS). *J Bone Miner Res*, 27, 2170–2178.
- Moayyeri, A., Hammond, C. J., Valdes, A. M., & Spector, T. D. (2012b). Cohort profile: TwinsUK and Healthy Ageing Twin Study. *International Journal of Epidemiology*. Available online, doi: 10.1093/ije/dyr207.
- Newton-Cheh, C., Johnson, T., Gateva, V., Tobin, M. D., Bochud, M., Coin, L., & Munroe, P. B. (2009). Genome-wide association study identifies eight loci associated with blood pressure. *Nature Genetics*, 41, 666–676.
- Nica, A. C., Parts, L., Glass, D., Nisbet, J., Barrett, A., Sekowska, M., . . . Spector, T. D., & MuTHER Consortium. (2011). The architecture of gene regulatory variation across multiple human tissues: the MuTHER study. *PLoS Genetics*, 7, e1002003.
- Nolte, I. M., Wallace, C., Newhouse, S. J., Waggott, D., Fu, J., Soranzo, N., & Bradbury, L. (2009). Common genetic variation near the phospholamban gene is associated with cardiac repolarisation: Meta-analysis of three genome-wide association studies. *PLoS One*, 4, e6138.
- Padmanabhan, S., Melander, O., Johnson, T., Di Blasio, A. M., Lee, W. K., Gentilini, D., & Dominiczak, A. F. (2010). Genome-wide association study of blood pressure extremes identifies variant near UMOD associated with hypertension. *PLoS Genetics*, 6, e1001177.
- Panoutsopoulou, K., Southam, L., Elliott, K. S., Wrayner, N., Zhai, G., Beazley, C., & Loughlin, J. (2011). Insights into the genetic architecture of osteoarthritis from stage 1 of the arcOGEN study. *Annals of the Rheumatic Diseases*, 70, 864–867.
- Repapi, E., Sayers, I., Wain, L. V., Burton, P. R., Johnson, T., Obeidat, M., & Tobin, M. D. (2010). Genome-wide association study identifies five loci associated with lung function. *Nature Genetics*, 42, 36–44.
- Richards, J. B., Kawoura, F. K., Rivadeneira, F., Stykarska, U., Estrada, K., Halldorsson, B. V., . . . Spector, T. D., for the GEFOS Consortium. (2009a). Collaborative meta-analysis: associations of 150 candidate genes with osteoporosis and osteoporotic fracture. *Annals of Internal Medicine*, 151, 528–537.
- Richards, J. B., Waterworth, D., O’Rahilly, S., Hivert, M.-F., Loos, R. J. F., Perry, J. R. B., . . . Spector, T. D., & GIANT Consortium. (2009b). A genome-wide association study reveals variants in ARL15 that influence adiponectin levels. *PLoS Genetics*, 5, e1000768.
- Smith, N. L., Chen, M. H., Dehghan, A., Strachan, D. P., Basu, S., Soranzo, N., & O’Donnell, C. J. (2010). Novel associations of multiple genetic loci with plasma levels of factor VII, factor VIII, and von Willebrand factor: The CHARGE (Cohorts for Heart and Aging Research in Genome Epidemiology) Consortium. *Circulation*, 121, 1382–1392.
- Spector, T. D., & Williams, F. M. (2006). The UK Adult Twin Registry (TwinsUK). *Twin Research & Human Genetics*, 9, 899–906.
- Speliotes, E. K., Willer, C. J., Berndt, S. I., et al. (2010). Association analyses of 249,796 individuals reveal 18 new loci associated with body mass index. *Nature Genetics*, 42, 937–948.
- Suhre, K., Shin, S. Y., Petersen, A. K., Monda, K. L., Thorleifsson, G., Jackson, A. U., & Loos, R. J. (2011). Human metabolic individuality in biomedical and pharmaceutical research. *Nature*, 477, 54–60.
- Willer, C. J., Speliotes, E. K., Loos, R. J., Li, S., Lindgren, C. M., Heid, I. M., . . . Hirschhorn, J. N., & Genetic Investigation of Anthropometric Traits Consortium. (2009). Six new loci associated with body mass index highlight a neuronal influence on body weight regulation. *Nature Genetics*, 41, 25–34.
- Zhai, G., Teumer, A., Stolk, L., Perry, J. R., Vandeput, L., Coviello, A. D., & Wallaschofski, H. (2011). Eight common genetic variants associated with serum DHEAS levels suggest a key role in ageing mechanisms. *PLoS Genetics*, 7, e1002025.
- Zhai, G., van Meurs, J. B., Livshits, G., Meulenbelt, I., Valdes, A. M., Soranzo, N., & Spector, T. D. (2009). A genome-wide association study suggests that a locus within the ataxin 2 binding protein 1 gene is associated with hand osteoarthritis: the Treat-OA consortium. *Journal of Medical Genetics*, 46, 614–616.
- Zhai, G., Wang-Sattler, R., Hart, D. J., Arden, N. K., Hakim, A. J., Illig, T., & Spector, T. D. (2010). Serum branched-chain amino acid to histidine ratio: A novel metabolomic biomarker of knee osteoarthritis. *Annals of the Rheumatic Diseases*, 69, 1227–1231.